Gyr A Monclonal Antibody (7F11) (N-terminal domain of Gyr A)



Product Description (Product Numbers #7F001, 7F005 and 7F100)

Gyr A antibody 7F11 is an IgG_{2b} monoclonal produced from tissue culture supernatant. It is purified on a Protein A column (Ey *et al.*, 1978) using the low-salt method. The antibody is supplied at a concentration requiring ~1:1000 dilution for Western blotting. **For in vitro laboratory research use only.**

Storage Buffer

Phosphate Buffered Saline (PBS) pH 7.2 Dilutions should be performed in PBS. Store at -20°C, (stable for at least 12 months undiluted).

Quality Control

The antibody is purified to >95% purity as judged by SDS-polyacrylamide gel electrophoresis.

Endonuclease assay: $0.5\mu g$ Supercoiled pBR322 incubated with the antibody at a concentration range of $0.01\mu g/ml-100\mu g/ml$ for 1 hour at $37^{\circ}C$ in the presence of 1 mM ATP, shows no detectable conversion of superhelical DNA to either open circular or linear forms when assayed by agarose gel electrophoresis.

Exonuclease assay: 1μg linear pBR322 was incubated with the antibody at a range of 0.01μg/ml-10μg/ml for 30 minutes at 37°C before being religated and transformed into competent JM109 cells. There was no detectable reduction in transformation efficiency.

References

Ey, P.L., S.J., & Jenkin, C.R. (1978). Isolation of pure IgG₁, IgG_{2a}, and IgG_{2b} immunoglobulins from mouse serum using Protein A sepharose. *Immunochemistry*. **15**: 429-436

Harlow, E. & Lane, D (1988). *Antibodies: a laboratory manual*. Cold Spring Harbor Laboratory Press. Cold Spring Harbor