

A. baumannii Gyrase Supercoiling Assay Kit



Product Description (Product Numbers ABS001 and ABS002)

A.baumannii gyrase is purified after heterologous expression in *E. coli* and is supplied as an equimolar complex of the subunits. The enzyme is supplied at a minimum concentration of 10 U/μl in Dilution Buffer. Store at -80°C. It is recommended that the enzyme is aliquoted to avoid repeated freeze-thaw cycles.

For *in vitro* laboratory research use only.

Dilution Buffer


50 mM KH₂PO₄
2 mM dithiothreitol
2 mM EDTA
30 % (w/v) glycerol

Assay Buffer (supplied as 5X stock)

35 mM Tris.HCl (pH 7.5)
24 mM KCl
4 mM MgCl₂
2 mM DTT
1.8 mM spermidine
1 mM ATP
6.5 % (w/v) glycerol
0.1 mg/ml albumin

Supercoiling Assay

1 U of gyrase is incubated with 0.5 μg of relaxed pBR322 in a reaction volume of 30 μl at 37 °C for 30 minutes in Assay Buffer. Gels are run in the absence of ethidium bromide or chloroquine.



Quality Control

Purity: The A and B subunits are purified to > 95 % purity as judged by SDS-polyacrylamide gel electrophoresis.

Endonuclease assay: 0.5 μg relaxed pBR322 incubated with 0.5 U of DNA gyrase for 1 hour at 37 °C in the presence of 1 mM ATP shows no detectable conversion of supercoiled DNA to either open circular or linear forms when assayed by agarose gel electrophoresis.

Reference

Hallett, P., Grimshaw, A.J., Wigley, D.B. and Maxwell, A. (1990). Cloning of the DNA gyrase genes under *tac* promoter control: overproduction of the gyrase A and B proteins. *Gene* 93: 139-142