

Staphylococcus aureus Topoisomerase IV



Product Description (Product Numbers SAT4001 and SAT4002)

S. aureus topoisomerase IV is purified after heterologous expression in *E. coli* and is supplied as a complex of the subunits. The enzyme is supplied at a minimum concentration of 10 U/ μ l in Dilution Buffer.

Store at -80°C. It is recommended that the enzyme is aliquoted to avoid repeated freeze-thaw cycles.

For *in vitro* laboratory research use only.

Dilution Buffer (1X)

50 mM Tris.HCl (pH 7.5)
1 mM DTT
1 mM EDTA
40 % (w/v) glycerol

Assay Buffer (supplied as 5X stock)

50 mM Tris.HCl (7.5)
5 mM MgCl₂
5 mM DTT
1.5 mM ATP
350 mM potassium glutamate
0.05 mg/ml albumin

DNA Storage Buffer

10 mM Tris.HCl (pH 7.5)
1 mM EDTA


Relaxation Assay

1 U of topoisomerase IV will relax 0.5 μ g of supercoiled pBR322 when incubated in 1X buffer in a total reaction volume of 30 μ l at 37°C for 30 minutes.

Gels are run in the absence of Ethidium Bromide or Chloroquine.

Sa topo IV

- +



Quality Control

- 1) Purity: *S. aureus* topoisomerase IV is purified to > 95 % purity as judged by SDS-polyacrylamide gel electrophoresis.
- 2) pBR322 was also incubated for 4 hrs in assay buffer + 10 mM MgCl₂ at 37 °C. These tests were negative for the formation of linear products, indicating the absence of nuclease contamination.
- 3) No activity was detectable when the single subunits were assayed alone